

## Product Information

**CAMK2δ, active, GST-tagged, human PRECISIO® Kinase recombinant, expressed in Sf9 cells**

Catalog Number **C0874**  
Lot Number 061M0737  
Storage Temperature  $-70^{\circ}\text{C}$

Synonyms: CAMKD, MGC44911

### Product Description

CAMK2δ is a serine/threonine protein kinase that is a member of the type II multifunctional  $\text{Ca}^{2+}$ /calmodulin-dependent protein kinase family. CAMK2δ is abundantly present in human cardiac and skeletal muscle, and its levels are increased in the hearts of patients suffering from cardiomyopathy.<sup>1</sup> In cardiomyocytes, stimulation of beta-1-adrenergic receptors leads to induction of apoptosis, an effect that is mediated by activation of CAMK2δ in a PKA-independent manner. In addition, expression studies have revealed the down regulation of CAMK2δ in human tumor cells.

This recombinant product was expressed by baculovirus in Sf9 insect cells using an N-terminal GST-tag. The gene accession number is NM 172115. It is supplied in 50 mM Tris-HCl, pH 7.5, with 150 mM NaCl, 0.25 mM DTT, 0.1 mM EGTA, 0.1 mM EDTA, 0.1 mM PMSF, and 25% glycerol.

Molecular mass: ~77 kDa

Purity:  $\geq 70\%$  (SDS-PAGE, see Figure 1)

Specific Activity: 4,981–6,739 nmole/min/mg  
(see Figure 2)


### Precautions and Disclaimer

This product is for R&D use only, not for drug, household, or other uses. Please consult the Material Safety Data Sheet for information regarding hazards and safe handling practices.


### Storage/Stability

The product ships on dry ice and storage at  $-70^{\circ}\text{C}$  is recommended. After opening, aliquot into smaller quantities and store at  $-70^{\circ}\text{C}$ . Avoid repeated handling and multiple freeze/thaw cycles.

**Figure 1.**  
SDS-PAGE Gel of Lot Number 061M0737:  
>85% (densitometry)



**Figure 2.**  
Specific Activity of Lot Number 061M0737:  
5,860 nmole/min/mg



### Procedure

#### Preparation Instructions

Kinase Assay Buffer – 25 mM MOPS, pH 7.2, 12.5 mM glycerol 2-phosphate, 25 mM  $\text{MgCl}_2$ , 5 mM EGTA, and 2 mM EDTA. Just prior to use, add DTT to a final concentration of 0.25 mM.

Kinase Dilution Buffer – Dilute the Kinase Assay Buffer 5-fold with a 50 ng/μl BSA solution.

Kinase Solution – Dilute the active CAMK2 $\delta$  (0.1  $\mu$ g/ $\mu$ l) with Kinase Dilution Buffer to the desired concentration. **Note:** The lot-specific specific activity plot may be used as a guideline (see Figure 2). It is recommended the researcher perform a serial dilution of active CAMK2 $\delta$  kinase for optimal results.

10 mM ATP Stock Solution – Dissolve 55 mg of ATP in 10 ml of Kinase Assay Buffer. Store in 200  $\mu$ l aliquots at –20 °C.

$\gamma$ -<sup>32</sup>P-ATP Assay Cocktail (250  $\mu$ M) – Combine 5.75 ml of Kinase Assay Buffer, 150  $\mu$ l of 10 mM ATP Stock Solution, 100  $\mu$ l of  $\gamma$ -<sup>32</sup>P-ATP (1 mCi/100  $\mu$ l). Store in 1 ml aliquots at –20 °C.

Substrate Solution – Dissolve the Autocamtide 2 synthetic peptide substrate (KKALRRQETVDAL-amide) in water at a final concentration of 1 mg/ml.

1% phosphoric acid solution – Dilute 10 ml of concentrated phosphoric acid to a final volume of 1 L with water.

#### Kinase Assay

This assay involves the use of the <sup>32</sup>P radioisotope. All institutional guidelines regarding the use of radioisotopes should be followed.

1. Thaw the active CAMK2 $\delta$ , Kinase Assay Buffer, Substrate Solution, and Kinase Dilution Buffer on ice. The  $\gamma$ -<sup>32</sup>P-ATP Assay Cocktail may be thawed at room temperature.
2. In a pre-cooled microcentrifuge tube, add the following solutions to a volume of 20  $\mu$ l:
  - 10  $\mu$ l of Kinase Solution
  - 7.5  $\mu$ l of Substrate Solution
  - 2.5  $\mu$ l of 5 mM CaCl<sub>2</sub> solution containing 0.75  $\mu$ g Calmodulin
3. Set up a blank control as outlined in step 2, substituting 7.5  $\mu$ l of cold water (4 °C) for the Substrate Solution.
4. Initiate each reaction with the addition of 5  $\mu$ l of the  $\gamma$ -<sup>32</sup>P-ATP Assay Cocktail, bringing the final reaction volume to 25  $\mu$ l. Incubate the mixture in a water bath at 30 °C for 15 minutes.
5. After the 15 minute incubation, stop the reaction by spotting 20  $\mu$ l of the reaction mixture onto an individually pre-cut strip of phosphocellulose P81 paper.

6. Air dry the pre-cut P81 strip and sequentially wash in the 1% phosphoric acid solution with constant gentle stirring. It is recommended the strips be washed a total of 3 times of ~10 minutes each.
7. Set up a radioactive control to measure the total  $\gamma$ -<sup>32</sup>P-ATP counts introduced into the reaction. Spot 5  $\mu$ l of the  $\gamma$ -<sup>32</sup>P-ATP Assay Cocktail on a pre-cut P81 strip. Dry the sample for 2 minutes and read the counts. Do not wash this sample.
8. Count the radioactivity on the P81 paper in the presence of scintillation fluid in a scintillation counter.
9. Determine the corrected cpm by subtracting the blank control value (see step 3) from each sample and calculate the kinase specific activity

#### Calculations:

1. Specific Radioactivity (SR) of ATP (cpm/nmole)

$$SR = \frac{\text{cpm of 5 } \mu\text{l of } \gamma\text{-}^{32}\text{P-ATP Assay Cocktail}}{\text{nmole of ATP}}$$

cpm – value from control (step 7)

nmole – 1.25 nmole (5  $\mu$ l of 250  $\mu$ M ATP Assay Cocktail)

2. Specific Kinase Activity (SA) (nmole/min/mg)

$$\text{nmole/min/mg} = \frac{\Delta\text{cpm} \times (25/20)}{SR \times E \times T}$$

SR = specific radioactivity of the ATP (cpm/nmole ATP)

$\Delta$ cpm = cpm of the sample – cpm of the blank (step 3)

25 = total reaction volume

20 = spot volume

T = reaction time (minutes)

E = amount of enzyme (mg)

#### References

1. Hoch, B. et al., Circ. Res., **84**, 713-721 (1999).
2. Zhu, W.-Z. et al., J. Clin. Invest., **111**, 617-625 (2003).

PRECISIO is a registered trademark of Sigma-Aldrich® Biotechnology LP and Sigma-Aldrich Co.

TD,MAM 06/11-1

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.