# Rev A\ 2013-12-05 \ SF-1313 DSCA\ VK

# ABL2 Hu-Cy3 SmartFlare™ RNA Detection Probe

Cat. # SF-1313

FOR RESEARCH USE ONLY

pack size: 50µL (250 rxns)

Store at 2-8℃, after reconstitution store at 23-27℃ DO NOT FREEZE



## **Product Data Sheet**

page 1

Accession #NM\_001136000.2

NOT FOR USE IN DIAGNOSTIC PROCEDURES

NOT FOR HUMAN OR ANIMAL CONSUMPTION

Species Hu

Gene Aliases: ARG; ABLL

### Confirmation of ABL2 SmartFlare Performance:

ABL2 SmartFlare probe has been tested in a buffer system to detect the release of the fluorophore in the presence of a complementary base pair sequence for each lot to confirm target specificity.

ABL2 SmartFlare probe has also been tested in a cell model system and demonstrated increased fluorescence in cells expressing the target compared to a scrambled negative control SmartFlare (Figure 1).

Mean Fluorescence Intensity (MFI) Values				
Unflared	Scrambled	ABL2		
39.25999999999998	270.48000000000002	1026.77		

**Figure 1:** ABL2 Mean Fluorescence Intensity (green) by flow cytometry in living HUVEC cells demonstrated a significant increase over unflared cells (blue) as well as scramble control (red). Data shown is representative.

### Storage and Handling:

Material has been 0.22µm filtered. Stable for 5 years at 2-8℃ degrees in lyophilized format ONLY. Room temperature is required for reconstituted product.

Warning-after reconstitution product is sensitive to cold and hot temperatures, a stable room temperature of 23-27°C is recommended.

### **Handling Recommendations:**

Reconstitute with sterile nuclease free water in a drop wise fashion tap tube repeatedly to fully dissolve lyophilized material.

Upon reconstitution, store at room temperature for up to 1 year protected from light. Product must be handled with gloves as product can be absorbed through the skin.

### **Recommended Cell Testing Protocol:**

(example: 30,000 cells in a 200µL media volume within each well of a 96 well plate)

- Reconstitute reagent in 50µL of sterile nuclease free water.
- Create a working solution based on your experiment by diluting 1:20 in sterile PBS
- Add 4µL directly to cells (at approx 80% confluency)
- Allow to incubate overnight for 16 hrs
- Detect using fluorescence detection platform of choice

